Unspoken Distress: Supporting Head and Neck Cancer Caregivers Through Early Psychosocial Intervention

Akiko Miyake, LCSW
NYU Langone Health Perlmutter Cancer Center

Social Work Department
Perlmutter Cancer Center

PURPOSE

To highlight the unique sources of distress experienced by caregivers of patients with head and neck cancer **(HNC)** and present an early oncology social work intervention model to improve preparedness, coping, and connection to support.

BACKGROUND/SIGNIFICANCE

Caregivers of patients with head and neck cancer often assume **complex clinical, emotional, and practical responsibilities** early in treatment. These demands can contribute to various levels of emotional distress. Distress may remain hidden because caregivers often prioritize the patient's needs, minimize their own distress, or feel shame about reactions such as resentment, fear, or loss of connection.

Early psychosocial intervention creates an opportunity to identify caregiver needs before distress intensifies and before treatment-related speech, swallowing, and appearance changes make communication and emotional connection more difficult.

WHY HEAD & NECK CANCER CAREGIVING IS UNIQUE

- Involves **complex, intimate, and highly visible care needs beyond traditional emotional support.**
- Caregivers may manage **communication changes**, **swallowing difficulties, modified diets, PEG feedings, and weight concerns**.
- Hands-on responsibilities can include **tracheostomy care, wound care, suctioning, pain management, and symptom monitoring**.
- Treatment-related **disfigurement and functional changes** may affect intimacy, social connection, family roles, and emotional adjustment.
- Because caregivers often prioritize the patient's needs, their own distress may remain **unspoken**, contributing to **fatigue, isolation, guilt, resentment**, and **self-neglect**.

FIGURE 1. UNIQUE ASPECTS OF HNC CAREGIVING



CLINICAL APPROACH / INTERVENTION MODEL

An oncology social work practice model was developed to engage caregivers at diagnosis and sustain support throughout treatment. The model emphasizes early identification of distress, interdisciplinary collaboration, and caregiver-centered education.

Assess → Prepare → Teach → Connect → Support

1. **Assess** caregiver distress, coping, and practical barriers
2. **Prepare** caregivers for role expectations and treatment demands
3. **Teach** brief coping skills, grounding and stress management
4. **Connect** caregivers to practical and psychosocial resources
5. **Support** emotional expression without judgment

CLINICAL OBSERVATIONS

Early psychosocial engagement with HNC caregivers suggests several practice-based benefits:

- Caregivers may feel **more prepared** for the caregiving role when support is introduced early.
- Caregivers may feel **less isolated** when given space to discuss stress, guilt, and uncertainty.
- Brief coping strategies may help caregivers **manage distress** during complex treatment phases.
- Early support may improve **caregiver confidence** in knowing when and how to seek help.
- Supported caregivers may be better able to **communicate** with the care team and assist with treatment adherence.

PRACTICE IMPLICATIONS

- Screen caregiver distress **early** and **reassess** throughout treatment.
- Normalize **difficult emotions**, including guilt, resentment, fear, grief and isolation.
- Prepare caregivers for **communication, nutrition, medical care**, and **role changes** before distress escalates.
- Embed **caregiver-centered education** and support into routine HNC care.
- Strengthen **interdisciplinary collaboration** among social work, nursing, nutrition, speech-language pathology, and mental health providers.

CONCLUSION

Caregivers of patients with head and neck cancer carry complex responsibilities that often remain unspoken or unrecognized. Early oncology social work intervention can help identify caregiver distress, improve preparedness, strengthen coping, and integrate caregiver support into standard cancer care.

As a practice-based model, this approach has not yet been evaluated through formal outcome measures and may vary based on staffing, referral workflows, and interdisciplinary resources. Future work should include standardized caregiver distress screening and evaluation of caregiver preparedness, coping, isolation, and resource connection over time.

REFERENCES

Baluni, P., Singh, R., Bhutani, R., & Mishra, A. (2024). *The impact of emotional distress on suicidal ideation in head and neck cancer patients: A review. Clinical Oncology and Research, 4*(4), 100408. https://doi.org/10.1016/j.oor.2024.100562

Christodoulou, L., Parpottas, P., & Petkari, E. (2024). *Psychological interventions to enhance positive outcomes in adult cancer caregivers: A systematic review. Journal of Health Psychology, 29*(7), 747–769. https://doi.org/10.1177/13591053241236254

Van Hof, K. S., Hoesseini, A., Dorr, M. C., Verdonck-de Leeuw, I. M., Jansen, F., Leemans, C. R., Takes, R. P., Terhaard, C. H. J., Baatenburg de Jong, R. J., Sewnaik, A., & Offerman, M. P. J. (2022). *Caregiver burden, psychological distress, and quality of life among informal caregivers of patients with head and neck cancer: A longitudinal study. International Journal of Environmental Research and Public Health, 19*(23), 16304. https://doi.org/10.3390/ijerph192316304

Van Hof, K. S., Hoesseini, A., Dorr, M. C., Verdonck-de Leeuw, I. M., Jansen, F., Leemans, C. R., Takes, R. P., Terhaard, C. H. J., Baatenburg de Jong, R. J., Sewnaik, A., & Offerman, M. P. J. (2023). *Unmet supportive care needs among informal caregivers of patients with head and neck cancer in the first two years after diagnosis and treatment: A prospective cohort study. Supportive Care in Cancer, 31*(5), 262. https://doi.org/10.1007/s00520-023-07670-1

Wang, T., Mazanec, S. R., & Voss, J. G. (2021). *Needs of informal caregivers of patients with head and neck cancer: A systematic review. Oncology Nursing Forum, 48*(1), 11–29. https://doi.org/10.1188/21.ONF.11-29

Zeng, Q., Ling, D., Chen, W., Liao, Y., An, R., & He, L. (2023). *Family caregivers' experiences of caring for patients with head and neck cancer: A systematic review and metasynthesis of qualitative studies. Cancer Nursing, 46*(1), E41–E61. https://doi.org/10.1097/NCC.0000000000001096